

CONSTRAINING MODELS OF MODIFIED GRAVITY WITH THE DOUBLE PULSAR PSR J0737-3039A/B SYSTEM

LORENZO IORIO

Viale Unità di Italia 68, 70125, Bari (BA), Italy
e-mail: lorenzo.iorio@libero.it

MATTEO LUCA RUGGIERO

Dipartimento di Fisica, Politecnico di Torino, Corso Duca degli Abruzzi 23, Torino, Italy
INFN, Sezione di Torino, Via Pietro Giuria 1, Torino, Italy
e-mail: matteo.ruggiero@polito.it

In this paper we use $\Delta P = -1.772341 \pm 13.153788$ s between the phenomenologically determined orbital period P_b of the PSR J0737-3039A/B double pulsar system and the purely Keplerian period $P^{(0)} = 2\pi\sqrt{a^3/G(m_A + m_B)}$ calculated with the system's parameters, determined independently of the third Kepler law itself, in order to put constraints on some models of modified gravity ($f(R)$, Yukawa-like fifth force, MOND). The major source of error affecting ΔP is not the one in the phenomenologically measured period ($\delta P_b = 4 \times 10^{-6}$ s), but the systematic uncertainty $\delta P^{(0)}$ in the computed Keplerian one due to the relative semimajor axis a mainly caused, in turn, by the errors in the ratio $\mathcal{R}$ of the pulsars' masses and in $\sin i$. We get $|\kappa| \leq 0.8 \times 10^{-26} \text{ m}^{-2}$ for the parameter that in the $f(R)$ framework is a measure of the non linearity of the theory, $|\alpha| \leq 5.5 \times 10^{-4}$ for the fifth-force strength parameter (for $\lambda \approx a = 0.006$ AU). The effects predicted by the strong-acceleration regime of MOND are far too small to be constrained with some effectiveness today and in the future as well. In view of the continuous timing of such an important system, it might happen that in the near future it will be possible to obtain somewhat tighter constraints.

PACS: 04.50.Kd; 04.80.Cc; 97.80.-d; 97.60.Gb

Keywords: Modified theories of gravity; Experimental tests of gravitational theories; Binaries; Pulsars

1. Introduction

One of the six Keplerian orbital elements in terms of which it is possible to parameterize the orbital motion of a pulsar in a binary system is the mean anomaly $\mathcal{M}$ defined as $\mathcal{M} \equiv n(t - T_0)$, where n is the mean motion and T_0 is the time of periastron passage. The mean motion $n \equiv 2\pi/P_b$ is inversely proportional to the time elapsed between two consecutive crossings of the periastron, i.e. the anomalistic period P_b . In Newtonian mechanics, for two point-like bodies, n reduces to the usual Keplerian expression $n^{(0)} = \sqrt{GM/a^3}$, where a is the semi-major axis of the relative orbit of the pulsar with respect to the companion and $M \equiv m_p + m_c$

is the sum of their masses. In pulsar timing the period P_b is very accurately determined in a phenomenological, model-independent way, so that it accounts for all the dynamical features of the system, not only those coming from the Newtonian point-like terms, within the measurement precision.

In this paper we wish to look for phenomenologically determined deviations from the third Kepler law in the PSR J0737-3039A/B system¹, consisting of two radio-pulsars moving along a moderately eccentric, 2.4 hr orbit (see Table 1 for its relevant orbital parameters), in order to dynamically constrain some models of modified gravity ($f(R)$, Yukawa-like fifth force, MOND). We believe that the

Table 1. Relevant orbital parameters of the PSR J0737-3039A/B system². The projected semimajor axis is defined as $x = (a_{bc}/c) \sin i$, where a_{bc} is the barycentric semimajor axis, the phenomenologically determined post-Keplerian parameter s is equal to $\sin i$ in general relativity, and i is the angle between the plane of the sky, perpendicular to the line-of-sight, and the orbital plane. We have conservatively quoted the largest error in s reported in Ref. 2. The parameter e is the eccentricity. The orbital period is known with a precision of 4×10^{-6} s. The quoted values of m_A and m_B were obtained in Ref. 2 by using the general relativistic expression of the post-Keplerian A's periastron advance for the sum of the masses $M = 2.58708(16)$, and the ratio of the masses $\mathcal{R} \equiv m_A/m_B = 1.0714(11)$ phenomenologically determined from both the projected semimajor axes x_A and x_B . The rotational period $\mathcal{P}_A = 2\pi/\Omega_A$ of PSR J0737-3039A/B A amounts to 22 ms, while $\mathcal{P}_B = 2\pi/\Omega_B = 2.75$ s.

P_b (d)	x_A (s)	s	e	m_A ($M_\odot$)	m_B ($M_\odot$)
0.10225156248(5)	1.415032(1)	0.99974(39)	0.0877775(9)	1.3381(7)	1.2489(7)

analysis presented here has the merit of quantitatively determining how useful a laboratory such as PSR J0737-3039A/B and a method like the orbital period can be in testing alternative theories of gravity by precisely individuating the major sources of errors. Indeed, as many different laboratories and methods as possible should be used to extensively tests a fundamental interaction as gravity. Such an analysis may also yield insights concerning future obtainable improvements in view of the continuous timing of such a binary system.

2. Deviations from the third Kepler law

Looking for genuine deviations from the third Kepler law—meant as discrepancy ΔP between the phenomenologically determined orbital period P_b and the purely Keplerian one $P^{(0)}$ calculated with the values of Table 1—from the PSR J0737-3039A/B timing data is possible because all the parameters entering ΔP have been just measured independently of the third Kepler law itself. Indeed, the relative semimajor axis entering $P^{(0)}$

$$a = (1 + \mathcal{R}) \left(\frac{cx_A}{s} \right) \quad (1)$$

is built in terms of the ratio $\mathcal{R}$ of the masses, the projected semimajor axis x_A and $\sin i$; the projected semimajor axes x_A and x_B were phenomenologically determined

from the timing data, the phenomenologically estimated post-Keplerian parameter s can be identified with $\sin i$ in general relativity, the ratio $\mathcal{R}$ has been phenomenologically determined from the ratio of the projected semimajor axes coming from the quite general relation, valid to at least the first post-Newtonian order^{3,4}

$$a_{\text{bc}}^{(\text{A})} m_{\text{A}} = a_{\text{bc}}^{(\text{B})} m_{\text{B}}, \quad (2)$$

and the sum of the masses has been, in turn, derived from the A's periastron advance which is, at present, the best determined post-Keplerian parameter. The third Kepler law only enters the expression of the mass functions

$$\mathcal{F}_{\text{A/B}} = \frac{m_{\text{B/A}}^3 \sin^3 i}{(m_{\text{A}} + m_{\text{B}})^2}, \quad (3)$$

which, instead, have not been used in obtaining the parameters of PSR J0737-3039A/B.

Table 1 yields

$$\Delta P \equiv P_{\text{b}} - P^{(0)} = -1.772341 \pm 13.153788 \text{ s}. \quad (4)$$

In Eq. (4) we have evaluated

$$\delta(\Delta P) \leq \delta P_{\text{b}} + \delta P^{(0)}, \quad (5)$$

with

$$\delta P^{(0)} \leq \delta P^{(0)}|_a + \delta P^{(0)}|_M + \delta P^{(0)}|_G. \quad (6)$$

The terms in Eq. (6) are

$$\begin{cases} \delta P^{(0)}|_a \leq 3\pi \sqrt{\frac{a}{GM}} \delta a = 12.218572 \text{ s}, \\ \delta P^{(0)}|_M \leq \pi \sqrt{\frac{a^3}{GM^3}} \delta M = 0.273248 \text{ s}, \\ \delta P^{(0)}|_G \leq \pi \sqrt{\frac{a^3}{G^3 M}} \delta G = 0.661964 \text{ s}, \end{cases} \quad (7)$$

so that

$$\delta P^{(0)} \leq 13.153784 \text{ s}. \quad (8)$$

In Eq. (7) we used the values of Table 1 for δM and⁵ $\delta G = 0.0010 \times 10^{-11} \text{ kg}^{-1} \text{ m}^3 \text{ s}^{-2}$; the uncertainty in a , which is the most important source of error, has been evaluated as

$$\delta a \leq \delta a|_{\mathcal{R}} + \delta a|_s + \delta a|_{x_{\text{A}}}, \quad (9)$$

with

$$\begin{cases} \delta a|_{\mathcal{R}} \leq \left(\frac{c x_{\text{A}}}{s}\right) \delta \mathcal{R} = 466,758 \text{ m}, \\ \delta a|_s \leq (1 + \mathcal{R}) \left(\frac{c x_{\text{A}}}{s^2}\right) \delta s = 342,879 \text{ m}, \\ \delta a|_{x_{\text{A}}} \leq (1 + \mathcal{R}) \left(\frac{c}{s}\right) \delta x_{\text{A}} = 621 \text{ m}. \end{cases} \quad (10)$$

Thus,

$$\delta a \leq 810,259 \text{ m.} \quad (11)$$

It is interesting to note that $\mathcal{R}$ and s have a major impact on the overall uncertainty in a ; our estimate has to be considered as conservative because we adopted for δs the largest value quoted in Ref. 2.

The following considerations about the treatment of errors are in order.

- Realistic upper bounds of the errors have been obtained by purposely linearly summing the various sources of uncertainty because of the existing correlations among the estimated parameters of the PSR J0737-3039A/B system²
- The uncertainties of Table 1 and used throughout the paper are twice the parameter uncertainties given by the software used in Ref. 2; the authors of such a work believe that the real measurement uncertainties are actually somewhat smaller than those quoted. If so, also our result for ΔP would, in fact, be more accurate. By the way, in view of the continuous timing of the PSR J0737-3039A/B system it is likely that in the near future the precision reached in determining its parameters will allow to better constrain ΔP , although it is difficult to quantify the extent of such an improvement because of the impact of x_B via $\mathcal{R}$.

3. Applications

In this Section we will look for deviations from the third Kepler law induced by small radial accelerations A_r , in general non-constant, induced by some modified models of gravity ($f(R)$, Yukawa-like fifth force, MOND). The effects of such accelerations, which are to be retained quite smaller than the Newtonian monopole, should, in principle, be treated perturbatively with, e.g., the Gauss perturbing equations for the variations of the Keplerian orbital elements of a test body. Indeed, while the average Newtonian acceleration for the PSR J0737-3039A/B system amounts to

$$\langle A_N \rangle = \frac{GM}{a^2 \sqrt{1-e^2}} = 446 \text{ m s}^{-2}, \quad (12)$$

the $f(R)$ acceleration (see Section 3.1) is about three orders of magnitude smaller, the Yukawa-type term (see Section 3.2) is five orders of magnitude smaller, while the MONDian correction (see Section 3.3) is even much smaller.

However, in view of the likely broad spectrum of readers, many of whom may not be fully acquainted with the rather specialistic tools of celestial mechanics, we will simply work with

$$\dot{\phi}^2 r = \frac{GM}{r^2} + A_r, \quad (13)$$

where ϕ is an azimuth angle, by neglecting the orbital eccentricity since the small corrections due to it would not affect the constraints set by the main terms.

3.1. *Extended theories of gravity: “ $f(R)$ ”*

It has been suggested that cosmic speed-up can be explained by generalizing general relativity introducing in the gravitational action terms non linear in the scalar curvature R (see Refs. 6, 7 and references therein). These theories are the so called $f(R)$ theories of gravity (or “extended theories of gravity”), where the gravitational Lagrangian depends on an arbitrary analytic function f of the scalar curvature R . If, on the one hand, extended theories of gravity seem to be appealing in cosmology, on the other hand, there is the need of checking $f(R)$ predictions with Solar System tests which general relativity passed with flying colors⁸. Actually, the debate on the compatibility of these theories with Solar System tests is still open (see Ref. 9 and Refs. 10, 11 and references therein). Here we do not enter the debate, but simply suggest how, on the basis of the approach already outlined in Refs. 10, 11, the data from the double pulsar can be used, at least in principle, to constrain the allowable analytical forms of the $f(R)$. We work in the Palatini formalism (for an approach to extended theories of gravity in the metric formalism see, for instance, Ref. 12 and references therein). The field equations of $f(R)$ theories, in the Palatini formalism, which have the explicit form

$$f'(R)R_{(\mu\nu)}(\Gamma) - \frac{1}{2}f(R)g_{\mu\nu} = \kappa T_{\mu\nu}^{mat} \quad (14)$$

$$\nabla_{\alpha}^{\Gamma}[\sqrt{g}f'(R)g^{\mu\nu}] = 0 \quad (15)$$

admit the spherically symmetric vacuum exact solution:

$$ds^2 = - \left(1 - \frac{2GM}{c^2 r} + \frac{kr^2}{3} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{2GM}{c^2 r} + \frac{kr^2}{3} \right)} + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\varphi^2, \quad (16)$$

which is referred to as Schwarzschild-de Sitter metric (see Ref. 13 for details on the derivation of this solution). The parameters appearing in Eq. (16) are the mass M of the spherically symmetric source of the gravitational field and k , which is related to the solutions $R = c_i$ of the structural equation

$$f'(R)R - 2f(R) = 0. \quad (17)$$

controlling the solutions of Eq. (14). Namely, it is $k = c_i/4 = R/4$. Indeed, we may say that k is a measure of the non-linearity of the theory (if $f(R) = R$, Eq. (17) has the solution $R = 0 \rightarrow k = 0$). In general relativity, the Schwarzschild-de Sitter solution corresponds to a spherically symmetric solution of Einstein equations, with a “cosmological” term $\Lambda g_{\mu\nu}$, Λ being the cosmological constant. In practice, it is $\Lambda = -k$ in our notation. As we can see from Eq. (16), the modifications to the solutions of the field equations due to $f(R)$ theories are given by a term proportional to the Ricci scalar. In what follows, we write the gravitational potential in the form

$$U \equiv U_0 + U_{\kappa}, \quad (18)$$

6

with

$$\begin{cases} U_0 = -\frac{GM}{r}, \\ U_\kappa = -\kappa r^2 \end{cases} \quad (19)$$

where $\kappa = k/3$, and we consider $\Delta U \equiv U_\kappa$ as a perturbation. In doing so, we neglect the effect of spatial curvature. From the perturbing potential U_k in Eq. (19) we obtain an entirely radial perturbing acceleration

$$\mathbf{A} = -2\kappa \mathbf{r}. \quad (20)$$

We may write

$$P_b \equiv P^{(0)} + P^{(k)} \quad (21)$$

where the contribution $\Delta P \equiv P^{(k)}$, due to the non linearity of the Lagrangian, by applying Eq. (13), turns out to be

$$\Delta P = -\frac{2\pi\kappa}{n^{(0)3}}, \quad (22)$$

from which κ can be evaluated

$$\kappa = -\frac{\Delta P n^{(0)3}}{2\pi} \quad (23)$$

Consequently, we may give the following estimates

$$\delta\kappa \leq \frac{n^{(0)3}}{2\pi} \delta(\Delta P) + \frac{3|\Delta P|n^{(0)2}}{2\pi} \delta n^{(0)}, \quad (24)$$

and

$$\kappa = (1 \pm 7) \times 10^{-10} \text{ s}^{-2} \quad (25)$$

Equivalently, restoring physical units

$$\kappa = (0.1 \pm 0.8) \times 10^{-26} \text{ m}^{-2}, \quad (26)$$

or, rephrased in terms of the cosmological constant,

$$|\Lambda| = 3\kappa = (0.3 \pm 2.5) \times 10^{-26} \text{ m}^{-2} \quad (27)$$

The estimates of Eq. (27) can be used to set constraints on the analytical form of the $f(R)$, according to an approach described in Refs. 10, 11. For instance, if we consider the Lagrangian

$$f(R) = R - \frac{\mu^4}{R}. \quad (28)$$

which mimics cosmic acceleration (even if it has well known instabilities see, e.g., Ref. 14), thanks to the estimates on κ we may set a limit on the parameter μ . We get

$$|\mu| \leq 10^{-20} \text{ eV}. \quad (29)$$

We remark that the estimates of Eq. (27) are of the same order of magnitudes of those obtained in Refs. 10, 11, where data coming from Solar System observations were used. So, as we did there, we can say that also data coming from pulsars are hardly fit to constrain $f(R)$, since the value Eq. (29) is remarkably greater than estimate⁶ $\mu \simeq 10^{-33}$ eV, needed for $f(R)$ gravity to explain the acceleration of the Universe without requiring dark matter.

3.2. Yukawa-like fifth force

Modifications of the gravitational potential leading to a Yukawa-like term arise in different contexts, such as brane-world models, scalar-vector-tensor theories of gravity, string theory, study of cosmological defects (see, e.g., Refs. 15, 16 and references therein). In general, a Yukawa-like force can be obtained from the potential¹⁷

$$U \equiv U_0 + U_Y, \quad (30)$$

with

$$\begin{cases} U_0 = -\frac{GM}{r}, \\ U_Y = -\frac{GM}{r} [\alpha \exp(-\frac{r}{\lambda})], \end{cases} \quad (31)$$

where $|\alpha|$ is the strength of the Yukawa-like force and λ its range¹⁸. The parameters α, λ have been bounded thanks to many laboratory experiments and observations, at different scales (see Ref. 19 and references therein). Here we show how the strength $|\alpha|$ of the Yukawa-like force can be constrained at a scale $\lambda \approx a \approx 10^9$ m, which is the length scale of the double pulsar PSR J0737-3039A/B. The perturbing potential $\Delta U \equiv U_Y$ in Eq. (31) leads to the following expression for the radial component of the corresponding perturbing acceleration:

$$A_Y = -\frac{GM\alpha}{r^2} \left(1 + \frac{r}{\lambda}\right) \exp\left(-\frac{r}{\lambda}\right). \quad (32)$$

From Eq. (32) it is possible to obtain, in the circular orbit approximation,

$$P = P^{(0)} \left[1 - \frac{\alpha}{2} \left(1 + \frac{r}{\lambda}\right) \exp\left(-\frac{r}{\lambda}\right)\right]. \quad (33)$$

Hence, on writing

$$P_b \equiv P^{(0)} + P^{(Y)}, \quad (34)$$

from Eq. (13) we obtain for the contribution $\Delta P \equiv P^{(Y)}$ the following expression

$$\Delta P = -\frac{\pi\alpha}{n^{(0)}} \left(1 + \frac{r}{\lambda}\right) \exp\left(-\frac{r}{\lambda}\right). \quad (35)$$

By assuming $\lambda \approx r \approx a$, we get

$$\begin{cases} \alpha \approx -\frac{\Delta P n^{(0)} \exp(-1)}{2\pi} = 0.7 \times 10^{-4}, \\ \delta\alpha \leq \frac{n^{(0)} \exp(-1)}{2\pi} \delta(\Delta P) + \frac{|\Delta P| \exp(-1)}{2\pi} \delta n^{(0)} = 5.5 \times 10^{-4}. \end{cases} \quad (36)$$

These estimates for α , should be compared with those obtained at different scales^{20,21,22}. In particular, in Ref. 21 it is showed that $|\alpha| \leq 10^{-9}$ at a scale corresponding to that of the double pulsar. In other words, our results are not compelling, since they are orders of magnitude greater than the best estimates already available. This means that the length scale of the hypothesized Yukawa-like fifth force is much greater or much smaller than the length-scale of the double pulsar PSR J0737-3039A/B ($a = 0.006$ AU). The results obtained in Ref. 22 for the Solar System point towards the second hypothesis, i.e. $\lambda < 0.006$ AU.

3.3. MOND

MOND predicts that the gravitational acceleration $\mathbf{A}_g$ felt by a particle in the field of a distribution of mass is

$$\mathbf{A}_g = \frac{\mathbf{A}_N}{\mu\left(\frac{A_g}{A_0}\right)}, \quad (37)$$

where $\mathbf{A}_N$ is the Newtonian acceleration, A_0 is an acceleration scale which different, independent ensembles of observations set to²³ $A_0 = 1.2 \times 10^{-10} \text{ m s}^{-2}$, and $\mu(x)$ is an interpolating function which approximates 1 for $x \gg 1$, i.e. for accelerations larger than A_0 ; for $x \ll 1$ $\mu(x) = x$, so that in such a strongly MONDian regime $A_g \approx \sqrt{A_N A_0}$. For a quite general class of interpolating functions, $\mu(x)$ can be cast into the form²⁴

$$\mu(x) \approx 1 - k_0 \left(\frac{1}{x}\right)^q, \quad (38)$$

which yields a modified gravitational acceleration²⁶

$$\mathbf{A}_g \approx \mathbf{A}_N \left[1 + k_0 \left(\frac{A_0}{A_N}\right)^q \right]. \quad (39)$$

Note that the most commonly used expressions for $\mu(x)$, i.e.²⁴

$$\mu(x) = \frac{x}{\sqrt{1+x^2}}, \quad (40)$$

and²⁵

$$\mu(x) = \frac{x}{1+x}, \quad (41)$$

can be obtained from Eq. (39) for $k_0 = 1/2, q = 2$ and $k_0 = 1, q = 1$, respectively. A useful review dealing, among other things, with many attempts to theoretically justify MOND is²⁷. Again, we may write

$$P_b \equiv P^{(0)} + P^{(M)}, \quad (42)$$

and, from Eq. (13) with A_r given by Eq. (39), on setting $e \rightarrow 0$, it is straightforward to obtain for the perturbation $\Delta P \equiv P^{(M)}$,

$$\Delta P = -\frac{P^{(0)} k_0}{2} \left(\frac{A_0}{A_N}\right)^q. \quad (43)$$

By defining

$$H_{k_0,q} \equiv -\frac{P^{(0)}k_0}{2} \left(\frac{A_0}{A_N} \right)^q, \quad (44)$$

it turns out that both Eq. (40) and Eq. (41) are compatible with the PSR J0737-3039A/B data: indeed,

$$\begin{cases} H_{1/2,2} = (-1.610 \pm 0.008) \times 10^{-22} \text{ s}, \\ H_{1,1} = (-1.431 \pm 0.004) \times 10^{-19} \text{ s}. \end{cases} \quad (45)$$

By the way, from Eq. (45) it is clear that the PSR J0737-3039A/B system is definitely unsuitable for testing MOND because discrepancies as small as those of Eq. (45) will never be measurable, falling well below even the precision in determining P_b . The same holds also for other binary systems, given the obtainable accuracy in determining their orbital periods. Solar System dynamics¹⁶ has proven to be more effective in constraining the interpolating function for $x \gg 1$.

4. Discussion and conclusions

In this paper we looked for deviations from the third Kepler law in the PSR J0737-3039A/B double pulsar system finding a discrepancy $\Delta P = -1.772341 \pm 13.153788$ s between the phenomenologically determined orbital period P_b and the purely Keplerian one $P^{(0)}$. While P_b is determined to a 10^{-6} s level, $P^{(0)}$ can be considered known only at $\approx 10^1$ s level after a conservative propagation of the uncertainties in the system's parameters entering its expression. The consistency of our analysis is assured by the fact that the values of the orbital parameters involved in the calculation of $P^{(0)}$ have been determined independently of the third Kepler law itself. The major source of error in $P^{(0)}$ is due to the semimajor axis which, in turn, is mainly affected by the ratio $\mathcal{R}$ of the pulsars' masses, which, in turn, depends on the poorly determined projected semimajor axis of B, and by $\sin i$. Continuous timing of the PSR J0737-3039A/B system might reduce such errors in the near future, perhaps yielding more precise results.

We used ΔP to constrain various theories of modified gravity like $f(R)$, Yukawa-type interactions and MOND. The bound on the parameter κ , that in $f(R)$ framework is a measure of the deviation of the theory from general relativity, is $|\kappa| \leq 0.8 \times 10^{-26} \text{ m}^{-2}$; differently speaking, the bound on κ can be seen as a bound on the cosmological constant $|\Lambda| = 3\kappa$. By assuming a range λ of the order of the PSR J0737-3039A/B system's size, we get $|\alpha| \leq 5 \times 10^{-4}$ for the strength of a putative Yukawa-like fifth force. In regard to MOND, the PSR J0737-3039A/B system is neatly unsuitable to test it because the order of magnitude of the predicted MONDian discrepancies is $\Delta P_{\text{MOND}} \approx 10^{-19} - 10^{-22}$ s.

Acknowledgements

L.I. gratefully thanks J. Katz for stimulating discussion. M.L.R. acknowledges financial support from the Italian Ministry of University and Research (MIUR) under the national program Cofin 2005 - *La pulsar doppia e oltre: verso una nuova era della ricerca sulle pulsar*. Both L.I. and M.L.R. gratefully acknowledge the referee for his/her useful comments which notably improved the manuscript.

References

1. M. Burgay *et al.*, *Nature* **426**, 531 (2003).
2. M. Kramer *et al.*, *Science* **314**, 97 (2006).
3. T. Damour and G. Schäfer, *Nuovo Cim.* **101**, 127 (1988).
4. T. Damour and J.H. Taylor, *Phys. Rev. D* **45**, 1840 (1992).
5. P.J. Mohr and B.N. Taylor, *Rev. Mod. Phys.* **77**, 1 (2005).
6. S.M. Carroll *et al.*, *Phys. Rev. D* **70**, 043528 (2004).
7. S. Nojiri and S.D. Odintsov, *Lectures given at 42nd Karpacz Winter School of Theoretical Physics*, hep-th/0601213.
8. C.M. Will, *Liv. Rev.* **9**, 3 (2006). (www.livingreviews.org/lrr-2006-3)
9. V. Faraoni, *Phys. Rev. D* **74**, 023529 (2006a).
10. M.L. Ruggiero and L. Iorio, *J. Cosmol. Astropart. Phys.* **0701**, 010 (2007).
11. G. Allemandi and M.L. Ruggiero, *Gen. Relativ. Gravit.* doi:10.1007/s10714-007-0441-3 (2007).
12. S. Capozziello, A. Stabile and A. Troisi, *Mod. Phys. Lett. A* **21**, 2291 (2006).
13. G. Allemandi *et al.*, *Gen. Relativ. Gravit.* **37**, 1891 (2005).
14. V. Faraoni, *Phys. Rev. D* **74**, 104017 (2006b).
15. E.G. Adelberger *et al.*, *Phys. Rev. Lett.* **98**, 131104 (2007).
16. M. Sereno and P. Jetzer, *Mon. Not. Roy. Astron. Soc.* **371**, 626 (2006).
17. O. Bertolami, J. Páramos and S.G. Turyshev, General Theory of Relativity: Will it survive the next decade?, in *Lasers, Clocks, and Drag-Free: Technologies for Future Exploration in Space and Tests of Gravity*, eds. H. Dittus, C. Lämmerzahl and S.G. Turyshev (Springer-Verlag, Berlin, 2006), p. 27.
18. E.G. Adelberger, B.R. Heckel and A.E. Nelson, *Ann. Rev. Nucl. Part. Sci.* **53**, 77 (2003).
19. O. Bertolami and F.M. Nunes, *Class. Quantum Grav.* **20**, L61 (2003).
20. O. Bertolami and J. Páramos, *Phys. Rev. D* **71**, 023521 (2005).
21. S. Reynaud and M.T. Jaekel, *Int. J. Mod. Phys. A* **20**, 2294 (2005).
22. L. Iorio, arXiv:0708.1080 [gr-qc] (2007).
23. R.H. Sanders and S.S. McGaugh, *Ann. Rev. Astron. Astrophys.* **40**, 263 (2002).
24. M. Milgrom, *Astroph. J.* **270**, 365 (1983).
25. B. Famaey and J. Binney, *Mon. Not. Roy. Astron. Soc.* **363**, 603 (2005).
26. C. Talmadge *et al.*, *Phys. Rev. Lett.* **61**, 1159 (1988).
27. J. Bekenstein, *Contemp. Phys.* **47**, 387 (2006).